

Bound State of Ultracold Fermions in the Spin Imbalanced Normal State

F. Fumarola, I.L. Aleiner and B.L. Altshuler

Physics Department, Columbia University, New York, NY 10027

(Dated: February 6, 2008)

We study a polarized Fermi gas and demonstrate that Fano-Feshbach (FF) resonances lead to the pairing of fermions and holes into long living massive bosonic modes (bifermions and biholes), which can be viewed as signatures of a first order quantum phase transition. These modes, which are neither conventional Cooper pairs nor closed channel bosonic dimers, have a peculiar dispersion relation and appear already in the weak coupling limit, i.e. for large FF detunings. Our result is in accord with claims of Schunck *et al.* [cond-mat/0702066] that by stabilizing the normal state the population imbalance still allows effect of s-wave fermionic pairing.

PACS numbers: 03.75.Ss, 05.30.Fk, 03.75.Kk

Quantum phase transitions in polarized Fermi gases have been a subject of investigation for many years [1–5]. In Ref. [1], it was demonstrated that spin polarization of electrons in a superconductor leads to a first order quantum phase transition. An applied magnetic field induces a Zeeman splitting E_Z in the energies of spin up and spin down electrons and imbalances the equilibrium densities of the two species. Eventually, for large enough E_Z the superconducting ground state becomes a polarized normal metal rather than a superconductor.

Let us define the imbalance in the system as $J = \frac{N_\uparrow - N_\downarrow}{N_\uparrow + N_\downarrow}$, $N_\uparrow$ and $N_\downarrow$ being the number of spin up and spin down fermions. To form an s-wave Cooper pair, two fermions have to be in different spin states. Therefore, for $J = 1$ the system has normal ground state. On the other hand, the ground state of a Fermi gas with no population imbalance ($J = 0$) is superfluid. Since no crossover between two states of different symmetry is possible, the system will remain normal even at zero temperature ($T = 0$) as long as J exceeds certain critical value J_c .

In superconductors, this translates into the existence of a critical magnetic field: in particular, the phase diagram in the presence of Zeeman (acting only on the spins) magnetic field [6] includes the BCS phase for $E_Z < \Delta$, a cohesistence region $\Delta < E_Z < \sqrt{2}\Delta$, and the normal state for $E_Z > \sqrt{2}\Delta$, where Δ is the BCS superconducting gap at zero temperature in the absence of Zeeman splitting. The elusive Fulde-Ferrell-Larkin-Ovchinnikov (FFLO) state [2, 3] appears in a narrow region below $E_Z = 1.508\Delta$: in this state, the mismatch of the spin up and spin down Fermi surfaces induces the electrons to form pairs with finite momentum. As a result, the superconducting gap becomes a periodic function of the space variables.

Experimental studies of superfluidity have been performed in ultracold Fermi gases [7, 8] in the BEC-BCS crossover, tuning the scattering length through Fano-Feshbach (FF) resonances [9, 10]. Effect of the polarization was achieved by creating imbalanced mixtures of ultracold ^6Li atoms in two different hyperfine states (which

we will refer to as spin states $\uparrow, \downarrow$). In the presence of the imbalance, anomalous density profiles [11] and the disappearance of quantum vortices [12] have been reported. While in metals orbital effects associated with the magnetic field can obscure the effects of Zeeman splitting, in neutral atomic gases this problem does not exist. Moreover, while in metals the relaxation due to spin orbit interaction tends to equilibrate the spin populations, in ultracold atomic gases the spin is a good quantum number, and the initial imbalance in the system is conserved. In these systems, the difference between the numbers $N_\uparrow$ and $N_\downarrow$ of majority and minority atoms can be encoded by the difference $2\hbar$ in the chemical potential of the two spin states, which plays the same role as the Zeeman splitting for electrons in metals.

In fact, the transition from superfluid to normal state in polarized Fermi gases (unlike the transition in an unpolarized gas) is an example of first order quantum phase transitions. Being discontinuous, first order quantum phase transitions appear to have no critical manifestations. However, as it was pointed out in [13], the signatures of such phase transitions are the anomalies (massive modes) in the *excitation* spectrum. These modes lead to sharp features in the observables and can determine the collective quantum dynamics. Moreover, the study of such modes may provide valuable information about the relaxation processes.

The identification of the massive modes and evaluation of their spectrum is a non-trivial model dependent task. The purpose of this Letter is to carry out this program for the normal state of resonantly coupled fermions with population imbalance. We prove that arbitrarily narrow FF resonances lead to the appearance of massive modes if J exceeds the critical value J_c so that the normal state is stable. We identify two kinds of bosonic quasiparticles: bifermions and biholes, having spin zero, associated with the pairing of fermions and holes. These modes have different features both from the closed channel dimers of the BEC-BCS crossover and from conventional Cooper pairs. They have a strongly J -dependent gap (shown in Fig. 1), and a peculiar dispersion relation, their energy




FIG. 1: Excitation gaps $\Omega_{bf}(q=0)$ and $\Omega_{bh}(q=0)$ of bifermions and biholes in units of Fermi energy as functions of detuning δ for fixed imbalance $J = \frac{N_{\uparrow} - N_{\downarrow}}{N_{\uparrow} + N_{\downarrow}} = 0.8$ (panel A) and as functions of the imbalance J for $\delta = 2.5\epsilon_F$ (panel B). The inset shows the behavior of the gaps for small imbalance, in the same units, together with the boundaries of the continuum (dashed lines). Here, the resonance width is $\gamma = 0.1$. The symmetry in the spectrum of bifermions and biholes ($\Omega_{bf} \sim -\Omega_{bh}$) is recovered for large detuning in panel A and for small imbalances in panel B.

being a decreasing function of momenta.

The FF resonant Fermi gas can be modeled by the following grandcanonical hamiltonian:

$$\begin{aligned} \hat{H} = & \int d\mathbf{r} \left[\sum_{\sigma} \hat{f}_{\sigma}^{\dagger} \epsilon_{\sigma}^f (-i\hbar\nabla) \hat{f}_{\sigma} + \hat{b}^{\dagger} \epsilon^b (-i\hbar\nabla) \hat{b} \right] \\ & + g \text{H.p} \int d\mathbf{r} \sum_{\sigma_1 \sigma_2} \left[\sigma_{\sigma_1 \sigma_2}^y \hat{b} \hat{f}_{\sigma_1}^{\dagger} \hat{f}_{\sigma_2}^{\dagger} \right], \quad (1) \\ \epsilon_{\sigma}^f = & \frac{\mathbf{p}^2}{2m} - \mu + 4\sigma h; \quad \epsilon^b = \frac{\mathbf{p}^2}{4m} - 2\mu + \delta_0, \end{aligned}$$

where σ^y is a Pauli matrix, H.p. denotes the Hermitian part of an operator, $\hat{f}_{\sigma}^{\dagger}(\mathbf{r})$, $[\hat{f}_{\sigma}(\mathbf{r})]$ create [annihilate] fermionic atoms with mass m and spin $\sigma = \pm 1/2$, while $\hat{b}^{\dagger}(\mathbf{r})$, $[\hat{b}(\mathbf{r})]$ create [annihilate] bosonic molecules; μ is the chemical potential. The bare detuning δ_0 depends on the high momentum cutoff $p_0 \rightarrow \infty$. It is related to the position of the FF resonance δ , which is independent of p_0 , and is connected with the s-wave scattering length a_s by:

$$\delta_0 = \delta + g^2 \int^{p_0} \frac{d^3 p}{(2\pi\hbar)^3} \frac{m}{p^2} \quad a_s = -\frac{mg^2}{4\pi\hbar\delta}. \quad (2)$$

The key dimensionless parameter is $\gamma = \frac{\sqrt{8}g^2 m^{3/2}}{4\pi^2 \hbar^3 \sqrt{\epsilon_F}}$, the ratio of the width of the FF resonance to the Fermi energy ϵ_F ($\epsilon_F \equiv \mu$ at $J = 0$). The model (1) suggests that in the weak coupling limit ($\gamma \ll 1$) that the system would be in the normal phase as long as $\delta > \delta_c(J) \sim 2\epsilon_F$ (Fig. 2). However, current experiments performed with wide




FIG. 2: (Color online) Schematic picture of the phase diagram at small coupling (for a complete discussion, see [5]). Arrows indicate the trajectories corresponding to the plots in Fig. 1. The FFLO phase lies in the red region at the superfluid/normal state boundary.

resonances ($\gamma \gg 1$) have shown that the system remains in the normal state down to much smaller values of the detuning: at resonance ($\delta = 0$), J_c was found as small as 0.7 [12]. We believe that our conclusions remain qualitatively valid for wide resonances and at resonance, as long as the normal state is stable.

In the case of narrow resonances ($\gamma \ll 1$), we can employ the one loop approximation for the bosonic Green function $\mathcal{D}(x_1, x_2)$, which amounts to the Dyson equation in Fig. 3, $\mathcal{D}^0$ and $\mathcal{G}^0$ being the non-interacting Green functions for the bosons and fermions correspondingly.

Since $\mathcal{D}$ is nothing but the two fermion scattering amplitude, bound states and resonances of the system are found by identifying its poles. Solving Eq. c) of Fig. 3 by means of Eqs. a) and b), after integration over frequencies, we find the equation for the dispersion relation $\Omega(q)$ of excitations in the particle-particle channel at zero temperature

$$\begin{aligned} \sum_{\pm} \int_0^{\infty} \sqrt{\epsilon} d\epsilon \int_{-1}^1 \frac{dx}{4} \left[-\frac{1}{2\epsilon} + \frac{\text{sgn}(\epsilon + \frac{q^2}{8m} - \mu \pm h + q\sqrt{\frac{\epsilon}{2m}}x)}{2\epsilon + \frac{q^2}{4m} - 2\mu - \Omega(q)} \right] \\ + \frac{\Omega(q) - \frac{q^2}{4m} + 2\mu - \delta}{\gamma\sqrt{\epsilon_F}} = 0. \quad (3) \end{aligned}$$

Here, h and the chemical potential μ are the Lagrange multipliers corresponding to the conserved quantities $N_{\uparrow} - N_{\downarrow}$ and $N_{\uparrow} + N_{\downarrow}$. In our discussion, h and μ can be calculated by setting $\gamma \rightarrow 0$, since any correction to the zero coupling limit would translate into corrections to Eq. (3) of higher order in the small parameter $\gamma \ll 1$. Hence, given the imbalance J and the density of atoms

n , one can determine h and μ from

$$\begin{aligned} \frac{\sqrt{2}}{3\pi^2} m^{3/2} (\mu + h)^{3/2} + \frac{\sqrt{2}}{3\pi^2} m^{3/2} (\mu - h)^{3/2} &= n \\ \frac{\sqrt{2}}{3\pi^2} m^{3/2} (\mu + h)^{3/2} - \frac{\sqrt{2}}{3\pi^2} m^{3/2} (\mu - h)^{3/2} &= Jn. \end{aligned} \quad (4)$$

Equation (3) has two solutions $\Omega_{bf}(q)$ and $\Omega_{bh}(q)$ which correspond to the two kinds of quasiparticles. These excitations have gaps, which can be found from the equation for Ω

$$\begin{aligned} \sqrt{\mu + \frac{\Omega}{2}} \left(\tanh^{-1} \left[\frac{\sqrt{\mu - h}}{\sqrt{\mu + \Omega/2}} \right] + \coth^{-1} \left[\frac{\sqrt{\mu + h}}{\sqrt{\mu + \Omega/2}} \right] \right) \\ - \sqrt{\mu + h} - \sqrt{\mu - h} + \frac{\Omega + 2\mu - \delta}{\gamma\sqrt{\epsilon_F}} = 0 \end{aligned} \quad (5)$$

The dependence of the gaps Ω_{bf} and Ω_{bh} on J and δ is plotted in Fig. 1.

Clearly, these quasiparticles have spin zero, and obey bosonic statistics. However, they are quite different from conventional Cooper pairs in that their energy is a decreasing function of momenta and their excitation gap strongly depends on the imbalance J of the system. Bifermions and biholes are also quite different from the closed channel bosonic dimers of the BEC-BCS crossover: in the limit $\delta \gg 2\epsilon_F$, such dimers are too energetic to play any role in the low energy physics of the Fermi gas. In contrast, bifermions and biholes can be observed experimentally even at large δ . Their excitation gap $\Omega_0 \sim 2h$ can be by orders of magnitude smaller than the binding energy δ of the bare bosonic dimers.

In the limit $J \ll 1$, the quasiparticle spectrum can be evaluated analytically. Equation (3) becomes

$$\sum_{\sigma, \tau = \pm 1} \frac{2\sigma h + v_F q + \sigma \tau \Omega(q)}{v_F q} \ln \left[\frac{2h + \sigma v_F q + \tau \Omega(q)}{\Delta} \right] = 4$$

with $h = \frac{2}{3} J \epsilon_F$. Here, Δ is the $T = 0$ superfluid order parameter, which satisfies the gap equation

$$\frac{\delta - 2\mu}{\gamma \epsilon_F} = \frac{1}{2} \int_{-\mu}^{\infty} d\xi \sqrt{1 + \frac{\xi}{\mu}} \left[\frac{1}{\sqrt{\xi^2 + \Delta^2}} - \frac{1}{\xi + \mu} \right] \quad (6)$$

For $\delta \gg 2\epsilon_F$, $\Delta \sim 8\epsilon_F \exp[-2 - \frac{\delta/\epsilon_F - 2}{\gamma}]$.

The dispersion relation of these quasiparticles is non-trivial. For $J \ll 1$, bifermions and biholes turn out to have opposite spectra. In this limit, as well as in the limit $\delta \gg \epsilon_F$ (see Fig. 1), bifermions and biholes can be viewed as particles and antiparticles. For $q \ll J\sqrt{m\epsilon_F}$,




FIG. 3: Diagrammatic representation of the Dyson equation for the particle-particle propagator, in the zero temperature formalism.

the dispersion relation is given by:

$$\begin{aligned} \Omega_{bf}(q) &= -\Omega_{bh}(q) = \Omega_{bf}(0) - \frac{1 + 2(\Omega_{bf}(0)/\Delta)^2}{6\Omega_{bf}(0)} v_F^2 q^2 \\ \Omega_{bf}(q=0) &= \sqrt{\left(\frac{4}{3} J \epsilon_F\right)^2 - \Delta^2} \end{aligned} \quad (7)$$

where $v_F = \sqrt{2\epsilon_F/m}$. According to Eq. (7), which is our main analytic result, the excitation energy is a decreasing function of momenta, and the excitation gap depends on the coupling between the fermions through the superfluid order parameter Δ .

In the one loop approximation, bifermions and biholes are undamped up to the critical value of momentum q_c [14], where their spectrum intersects a cut of the bosonic Green function (Fig. 4). For $q > q_c$, the collective modes develop a nonzero imaginary part.

The vanishing of $\Omega_{bf}(q=0)$ and $\Omega_{bh}(q=0)$ at $J = \frac{3}{4} \frac{\Delta}{\epsilon_F}$ manifests the instability of the normal state with respect to formation of the homogeneous Δ for $J < \frac{3}{4} \frac{\Delta}{\epsilon_F}$. This corresponds to the first order phase transition mentioned above. As to the instability toward inhomogeneous $\Delta(\mathbf{q}) \sim e^{i\mathbf{q}\cdot\mathbf{r}}$, it appears at $\tilde{J}_c = 1.13 \frac{\Delta}{\epsilon_F}$ [3], corresponding to a second order quantum phase transition to the FFLO phase with broken translational symmetry.

The critical behavior near the second order transition between the normal and FFLO state at $J < J_c$, as well as the collective excitations, definitely deserve theoretical analysis. Here we can note just that the bifermion and bihole modes have little to do with these problems. Namely, the analysis of the solution of Eq. (3) on the physical and nonphysical sheets suggests that the FFLO transition manifests itself in the appearance of new diffusive branch rather than in softening of the mode (7).




FIG. 4: (Color online) The main plot shows the dispersion relations of bifermions and biholes, for $J = J_c$ (frequencies are in units of $2h$, momenta in units of $2h/v_F$). The value q_c for which the spectrum intersects the cuts $\Omega = \pm(2h - v_F q)$ (red lines in the main plot) is plotted in the inset as a function of $J\epsilon_F/\Delta$, in units of Δ/v_F .

In the context of disordered superconductors, the existence of massive modes was discovered through measurements of the tunneling density of states [15], and the theory of the effect was constructed in Ref. [13]. We believe that bifermions and biholes in ultracold atomic gases should manifest themselves in measurements of radio-frequency spectroscopy, which have detected superfluidity at $J = 0$ [8]. Possible additional experimental applications of those modes will be discussed elsewhere [16].

In conclusion, we have shown that FF resonances lead to collective modes in a Fermi gas with spin imbalance. As long as the imbalance is large enough to prevent the formation of inhomogeneous configurations, the physics is dominated by the appearance of bosonic quasiparticles (bifermions and biholes). These quasiparticles have finite gap, anomalous dispersion relation, and exist as long as the system is in the normal state.

Added Note. After this paper was completed, experimental data were published by Shunck *et al.* [17], suggesting that s-wave pairing of atoms indeed takes place in the spin imbalanced normal phase of ultracold fermionic gases, as an explanation of anomalies in the population of imbalanced ^6Li atoms after applied radiofrequency. We find it likely that bifermion modes, which we discussed here provide a coherent interpretation of those anomalies.

- [1] A. M. Clogston, Phys. Rev. Lett. **9**, 266 (1962); B. S. Chandrasekhar, Appl. Phys. Lett. **1**, 7 (1962).
- [2] P. Fulde and R. A. Ferrell, Phys. Rev. **135**, A550 (1964).
- [3] A.I. Larkin and Y.N. Ovchinnikov, Zh. Eksp. Teor. Fiz. **47**, 1136 (1964) [Sov. Phys. JETP **20**, 762 (1965)].
- [4] G. Sarma, J. Phys. Chem. Solids **24**, 1029 (1963); R. Combescot, Europhys. Lett. **55**, 150 (2001); H. Caldas, Phys. Rev. A **69**, 063602 (2004); A. Sedrakian *et al.*, Phys. Rev. A **72**, 013613 (2005); U. Lombardo *et al.*, Phys. Rev. C **64**, 064314 (2001); D. T. Son and M. A. Stephanov, Phys. A **72**, 013614 (2006); L. He, M. Jin and P. Zhuang, Phys. B **73**, 214527 (2006); P. F. Bedaque, H. Caldas, and G. Rupak, Phys. Rev. Lett. **91**, 247002 (2003); C.-H. Pao, S.-T. Wu, and S.-K. Yip, Phys. Rev. B **73**, 132506 (2006); D. E. Sheehy and L. Radzihovsky, Phys. Rev. Lett. **96**, 060401 (2006); W. V. Liu and F. Wilczek, Phys. Rev. Lett. **90**, 047002 (2003); J. Carlson and S. Reddy, Phys. Rev. Lett. **95**, 060401 (2005); P. Castorina *et al.*, Phys. Rev. A **72**, 025601 (2005); T. Mizushima, K. Machida, and M. Ichioka, Phys. Rev. Lett. **94**, 060404 (2005).
- [5] D. E. Sheehy and L. Radzihovsky, cond-mat/0607803.
- [6] The orbital effect of the magnetic field can be made negligible in superconductors in the presence of magnetic impurities or in restricted geometries.
- [7] K. M. O'Hara *et al.*, Science **298**, 2179 (2002); C. A. Regal, Nature **424**, 47 (2003); M. Greiner, C. A. Regal, and D. S. Jin, Nature **426**, 537 (2003); S. Jochim *et al.*, Science **302**, 2101 (2003); M. W. Zwierlein *et al.*, Phys. Rev. Lett. **91**, 250401 (2003); C. A. Regal, M. Greiner and D. S. Jin, Phys. Rev. Lett. **92**, 040403 (2004); M. W. Zwierlein *et al.*, Phys. Rev. Lett. **92**, 120403 (2004); J. Kinast *et al.*, Phys. Rev. Lett. **92**, 150402 (2004); M. Bartenstein *et al.*, Phys. Rev. Lett. **92**, 203201 (2004); T. Bourdel *et al.*, Phys. Rev. Lett. **93**, 050401 (2004); M. W. Zwierlein *et al.*, Nature **435**, 1047-1051 (2005); G. B. Partridge *et al.*, Phys. Rev. Lett. **95**, 020404 (2005); J. Kinast *et al.*, Science **307**, 1296 (2005).
- [8] C. Chin *et al.*, Science **305**, 1128 (2004).
- [9] U. Fano, Nuovo Cimento, **12**, 156 (1955); Phys. Rev. A **124**, 1866 (1961); H. Feshbach, Ann. Phys. **5**, 357 (1958).
- [10] S. Inouye, *et al.*, Nature, **392**, 151 (1998); E. A. Donley *et al.*, Nature **417**, 529-533 (2002); E. Timmermans, *et al.*, Phys. Rep. **315**, 199 (1999); D. S. Petrov, C. Salomon and G. V. Shlyapnikov, Phys. Rev. Lett. **93**, 090404 (2004); A.V. Andreev, V. Gurarie and L. Radzihovsky, Phys. Rev. Lett. **93**, 130402 (2004); Q. Chen *et al.*, Phys. Rep. **412**, 1-88 (2005); D. S. Petrov, C. Salomon, and G. V. Shlyapnikov, J. Phys. B **38**, S645 (2005); V. Gurarie and L. Radzihovsky, cond-mat/0611022.
- [11] G. B. Partridge *et al.*, Science, **311**, 503 (2006).
- [12] M. W. Zwierlein *et al.*, Science, **311**, 492 (2006); Nature, **442**, 54 (2006).
- [13] I.L. Aleiner and B.L. Altshuler, Phys. Rev. Lett. **79**, 4242 (1997); Hae-Young Kee, I. L. Aleiner, and B. L. Altshuler, Phys. Rev. B **58**, 5757 (1998).
- [14] Higher loops corrections would provide a finite imaginary part to the spectrum of bifermions and biholes, because the decay into four fermions is not forbidden by momentum conservation even for $q = 0$.
- [15] W. Wu, J. Williams, and P.W. Adams, Phys. Rev. Lett. **77**, 1139 (1996); V. Yu. Butko, P. W. Adams and I. L.

- Aleiner, Phys. Rev. Lett. **82**, 4284 (1999).
- [16] F.Fumarola, I.L. Aleiner and B.L. Altshuler, in preparation.
- [17] C.H. Schunck *et al.*, cond-mat/0702066.